

# $\mathcal{B}(H)$ has a pure state that is not multiplicative on any masa

Charles Akemann<sup>1\*</sup> and Nik Weaver<sup>5</sup>

<sup>1</sup>Department of Mathematics, University of California, Santa Barbara, CA 93106; and <sup>5</sup>Department of Mathematics, Washington University in Saint Louis, Saint Louis, MO 63130

Edited by Richard V. Kadison, University of Pennsylvania, Philadelphia, PA, and approved February 7, 2008 (received for review February 5, 2008)

**Assuming the continuum hypothesis, we prove that  $\mathcal{B}(H)$  has a pure state whose restriction to any masa is not pure. This resolves negatively old conjectures of Kadison and Singer and of Anderson.**

bounded operators | Hilbert space

Let  $H$  be a separable infinite-dimensional Hilbert space and let  $\mathcal{B}(H)$  be the algebra of bounded operators on  $H$ . Kadison and Singer (1) suggested that every pure state on  $\mathcal{B}(H)$  would restrict to a pure state on some maximal abelian self-adjoint subalgebra (masa). Anderson (2) formulated the stronger conjecture that every pure state on  $\mathcal{B}(H)$  is *diagonalizable*, that is, of the form  $f(A) = \lim_U \langle Ae_n, e_n \rangle$  for some orthonormal basis  $(e_n)$  and some ultrafilter  $U$  over  $\mathbb{N}$ .

An *atomic masa* is the set of all operators that are diagonalized with respect to some given orthonormal basis of  $H$ . Anderson's conjecture is related to a fundamental problem in  $C^*$ -algebras also raised in ref. 1 and now known as the Kadison–Singer problem, which asks whether every pure state on an atomic masa of  $\mathcal{B}(H)$  has a unique extension to a pure state on  $\mathcal{B}(H)$ . If  $(e_n)$  is an orthonormal basis of  $H$ , then every pure state  $f_0$  on the corresponding atomic masa  $\mathcal{M}$  has the form  $f_0(A) = \lim_U \langle Ae_n, e_n \rangle$  for some ultrafilter  $U$  over  $\mathbb{N}$  and all  $A \in \mathcal{M}$ , and Anderson (3) showed that the same formula, now for  $A \in \mathcal{B}(H)$ , defines a pure state  $f$  on  $\mathcal{B}(H)$ . Thus, a positive solution to the Kadison–Singer problem would say that  $f$  is the only pure state on  $\mathcal{B}(H)$  that extends  $f_0$ .

In the presence of a positive solution to the Kadison–Singer problem, Anderson's conjecture is equivalent to the weaker statement that every pure state on  $\mathcal{B}(H)$  restricts to a pure state on some atomic masa. However, assuming the continuum hypothesis, we show that this weaker statement is false; in fact, there exist pure states on  $\mathcal{B}(H)$  whose restriction to any masa is not pure. It follows that there are pure states on  $\mathcal{B}(H)$  that are not diagonalizable. It seems likely that the statement “every pure state on  $\mathcal{B}(H)$  restricts to a pure state on some atomic masa” is also consistent with standard set theory. This, together with a positive solution to the Kadison–Singer problem, would imply the consistency of a positive answer to Anderson's conjecture.

The key lemma we need is the following. Let  $\mathcal{K}(H)$  be the algebra of compact operators on  $H$ , let  $\mathcal{C}(H) = \mathcal{B}(H)/\mathcal{K}(H)$  be the Calkin algebra, and let  $\pi: \mathcal{B}(H) \rightarrow \mathcal{C}(H)$  be the natural quotient map. We also write  $\dot{a}$  for  $\pi(a)$ , for any  $a \in \mathcal{B}(H)$ .

**Lemma 0.1.** *Let  $\mathcal{A}$  be a separable  $C^*$ -subalgebra of  $\mathcal{B}(H)$  which contains  $\mathcal{K}(H)$ , let  $f$  be a pure state on  $\mathcal{A}$  that annihilates  $\mathcal{K}(H)$ , and let  $\mathcal{M}$  be a masa of  $\mathcal{B}(H)$ . Then there is a pure state  $g$  on  $\mathcal{B}(H)$  that extends  $f$  and whose restriction to  $\mathcal{M}$  is not pure.*

**Proof:** By Proposition 6 of ref. 4 we can find an infinite-rank projection  $p \in \mathcal{B}(H)$  such that

$$p\dot{a}p = f(a)p \quad [1]$$

for all  $a \in \mathcal{A}$ .

Lemma 1.4 and Theorem 2.1 of ref. 5 imply that  $\pi(\mathcal{M})$  is a masa of  $\mathcal{C}(H)$ . It follows that there is a projection  $q \in \mathcal{M}$  such that  $\dot{q}$  neither contains nor is orthogonal to  $\dot{p}$ . Otherwise  $\dot{p}$  would be

in the commutant of  $\pi(\mathcal{M})$ , and hence would belong to  $\pi(\mathcal{M})$  by maximality. But this would mean  $\dot{p}$  is minimal in  $\pi(\mathcal{M})$  because any nonzero projection below  $\dot{p}$  neither contains nor is orthogonal to  $\dot{p}$ , and  $\pi(\mathcal{M})$  has no minimal projections.

Let  $\phi: \mathcal{C}(H) \rightarrow \mathcal{B}(K)$  be an irreducible representation of the Calkin algebra. It is faithful because  $\mathcal{C}(H)$  is simple. Therefore,  $\phi(\dot{q})$  neither contains nor is orthogonal to  $\phi(\dot{p})$ , so we can find a unit vector  $v \in K$  in the range of  $\phi(\dot{p})$  which is neither contained in nor orthogonal to the range of  $\phi(\dot{q})$ . Finally, define  $g(a) = \langle \phi(\dot{a})v, v \rangle$  for all  $a \in \mathcal{B}(H)$ . This is a pure state on  $\mathcal{B}(H)$  because  $\phi \circ \pi$  is an irreducible representation of  $\mathcal{B}(H)$ . It extends  $f$  because, using Eq. 1,

$$g(a) = \langle \phi(\dot{a})v, v \rangle = \langle \phi(\dot{a})\phi(\dot{p})v, \phi(\dot{p})v \rangle = \langle \phi(\dot{p}\dot{a}p)v, v \rangle = \langle f(a)\phi(\dot{p})v, v \rangle = f(a)$$

for all  $a \in \mathcal{A}$ . Finally, its restriction to  $\mathcal{M}$  is not pure because the projection  $q \in \mathcal{M}$  has the property that

$$g(q) = \langle \phi(\dot{q})v, v \rangle$$

is strictly between 0 and 1, since  $v$  is neither contained in nor orthogonal to the range of  $\phi(\dot{q})$ .  $\square$

**Theorem 0.2.** *Assume the continuum hypothesis. Then there is a pure state on  $\mathcal{B}(H)$  whose restriction to any masa is not pure.*

**Proof:** Let  $(a_\alpha)$ ,  $\alpha < \aleph_1$ , enumerate the elements of  $\mathcal{B}(H)$ . Since every von Neumann subalgebra of  $\mathcal{B}(H)$  is countably generated, a simple cardinality argument shows that there are only  $\aleph_1$  such subalgebras. Hence,  $\mathcal{B}(H)$  has only  $\aleph_1$  masas. Let  $(\mathcal{M}_\alpha)$ ,  $\alpha < \aleph_1$ , enumerate the masas of  $\mathcal{B}(H)$ .

We now inductively construct a nested transfinite sequence of unital separable  $C^*$ -subalgebras  $\mathcal{A}_\alpha$  of  $\mathcal{B}(H)$  together with pure states  $f_\alpha$  on  $\mathcal{A}_\alpha$  such that for all  $\alpha < \aleph_1$

1.  $a_\alpha \in \mathcal{A}_{\alpha+1}$ .
2. if  $\beta < \alpha$  then  $f_\alpha$  restricted to  $\mathcal{A}_\beta$  equals  $f_\beta$ .
3.  $\mathcal{A}_{\alpha+1}$  contains a projection  $q_\alpha \in \mathcal{M}_\alpha$  such that  $0 < f_{\alpha+1}(q_\alpha) < 1$ .

Begin by letting  $\mathcal{A}_0$  be any separable  $C^*$ -subalgebra of  $\mathcal{B}(H)$  that is unital and contains  $\mathcal{K}(H)$  and let  $f_0$  be any pure state on  $\mathcal{A}_0$  that annihilates  $\mathcal{K}(H)$ . At successor stages, use the lemma to find a projection  $q_\alpha \in \mathcal{M}_\alpha$  and a pure state  $g$  on  $\mathcal{B}(H)$  such that  $g|_{\mathcal{A}_\alpha} = f_\alpha$  and  $0 < g(q_\alpha) < 1$ . By Lemma 4 of ref. 6 there is a separable  $C^*$ -algebra  $\mathcal{A}_{\alpha+1} \subseteq \mathcal{B}(H)$  that contains  $\mathcal{A}_\alpha$ ,  $q_\alpha$ , and  $g$ , and such that the restriction  $f_{\alpha+1}$  of  $g$  to  $\mathcal{A}_{\alpha+1}$  is pure. To see this, write  $\mathcal{B}(H)$  as the union of a continuous nested transfinite sequence of separable  $C^*$ -algebras  $\mathcal{B}_\gamma$  such that  $\mathcal{B}_0$  is the  $C^*$ -algebra

Author contributions: C.A. and N.W. designed research, performed research, analyzed data, and wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

\*To whom correspondence should be addressed. E-mail: akemann@math.ucsb.edu.

© 2008 by The National Academy of Sciences of the USA

generated by  $\mathcal{A}_\alpha$ ,  $a_\alpha$ , and  $q_\alpha$ . The cited lemma guarantees that the restriction of  $g$  to some  $\mathcal{B}_\gamma$  will be pure. Thus, the construction may proceed. At limit ordinals  $\alpha$ , let  $\mathcal{A}_\alpha$  be the closure of  $\bigcup_{\beta < \alpha} \mathcal{A}_\beta$ . The state  $f_\alpha$  is determined by the condition  $f_\alpha|_{\mathcal{A}_\beta} = f_\beta$ , and it is easy to see that  $f_\alpha$  must be pure. [If  $g_1$  and  $g_2$  are states on  $\mathcal{A}_\alpha$  such that  $f_\alpha = (g_1 + g_2)/2$ , then for all  $\beta < \alpha$  purity of  $f_\beta$  implies that  $g_1$  and  $g_2$  agree when restricted to  $\mathcal{A}_\beta$ ; thus  $g_1 = g_2$ .] This completes the description of the construction.

Now define a state  $f$  on  $\mathcal{B}(H)$  by letting  $f|_{\mathcal{A}_\alpha} = f_\alpha$ . By the reasoning used immediately above,  $f$  is pure, and since  $0 < f(q_\alpha) < 1$  for all  $\alpha$ , the restriction of  $f$  to any masa is not pure.

It is interesting to contrast Theorem 0.2 with Theorem 9 of ref. 4, which states that (assuming the continuum hypothesis) any state on  $\mathcal{C}(H)$  restricts to a pure state on some masa of  $\mathcal{C}(H)$ . This does not conflict with our result because there are many masas of  $\mathcal{C}(H)$  that do not come from masas of  $\mathcal{B}(H)$  (regardless of the truth of the continuum hypothesis). Indeed,  $\mathcal{B}(H)$  has  $2^{\aleph_0}$  masas but  $\mathcal{C}(H)$  has  $2^{2^{\aleph_0}}$  masas. This can be seen by first finding  $2^{\aleph_0}$  mutually orthogonal nonzero projections  $p_\alpha$  in  $\mathcal{C}(H)$ , then finding projections  $q_\alpha^1, q_\alpha^2 < p_\alpha$  such that  $q_\alpha^1 q_\alpha^2 \neq q_\alpha^2 q_\alpha^1$  for each  $\alpha$ , and finally for each set  $S \subseteq 2^{\aleph_0}$  choosing a masa of  $\mathcal{C}(H)$  that contains  $\{q_\alpha^1 : \alpha \in S\}$  and  $\{q_\alpha^2 : \alpha \notin S\}$ . It is easy to see that this produces  $2^{2^{\aleph_0}}$  distinct masas.

1. Kadison RV, Singer IM (1959) Extensions of pure states. *Am J Math* 81:383–400.
2. Anderson J (1981) A conjecture concerning the pure states of  $\mathcal{B}(H)$  and a related theorem. *Topics in Modern Operator Theory* (Birkhäuser, Basel), pp 27–43.
3. Anderson J (1979) Extreme points in sets of positive linear maps on  $\mathcal{B}(H)$ . *J Funct Anal* 31:195–217.
4. Anderson J (1979) Pathology in the Calkin algebra. *J Operator Theory* 2:159–167.
5. Johnson BE, Parrott SK (1972) Operators commuting with a von Neumann algebra modulo the set of compact operators. *J Funct Anal* 11:39–61.
6. Akemann C, Weaver N (2004) Consistency of a counterexample to Naimark's problem. *Proc Natl Acad Sci USA* 101:7522–7525.